



Application of Improved Direct Calibration for Hyperspectral image processing : Detecting peanut traces in wheat flour



Puneet Mishra¹, Jean Michel Roger², Pilar Barreiro¹, Belén Diezma¹,
Ana Herrero-Langreo², Lourdes Lleó¹, Nathalie Gorretta²



¹LPF-TAGRALIA, UPM-CEI Moncloa. ETSI Agrónomos, Avda Complutense s/n. 28040, Madrid, Spain
²Irstea, UMR ITAP, 361 Rue J.F. Breton, 34196 Montpellier Cedex 5, France



Background



In current **industrial environments** there is an increasing **need** for practical and **in-line detection of foreign materials** in powder food processing lines. This demand is **especially** important regarding to the detection of **highly allergenic products**, such as **peanut**. Peanut are the leading cause of fatalities from food-induced allergic reactions², being avoidance the primary management of these allergies³. Adapted detection methods require:

- **High sensitivity**, to detect small traces
- **Robustness** to industrial environments

Objective

Evaluate the feasibility of hyperspectral (HS) imaging and Improved Direct Calibration (IDC¹) for the detection of peanut traces in wheat flour.

Materials

- Wheat flour (125-100 and 212-160 µm), “Coeur de Blé” from manufacturer MasterChef
- Peanut (500-1000 µm) : obtained from European Commission Institute for Reference Materials and Measurements (IRMM-481kit)
- Aluminum platforms (36 cm² and 95 cm²) (Fig. a.)
- Eleven samples were made: pure peanut, pure wheat flour, samples with known position of peanut on the surface and eight homogeneously mixed samples from 10% to 0.01% by weight.
- HySpex SWIR-320m-e (1000-2500 nm) line-scan push broom camera by Norsk Elektro Optikk, Norway



Fig a. Sample preparation

Methods

1. Loadings calculation

Based on:

- **Expert information:** Pure spectra from each product can be used for the loadings calculation, allowing **practical, specific and sensitive** product identification.

- **Experimental information:** “Toxic” information, such as variation around the mean can be removed to improve the **robustness** of the method.

1.1 Spectral pretreatment: SNV and Savitzky-Golay

1.2 Loadings calculation:

$$b_{IDC} = \Sigma_{IDC} k' (k \Sigma_{IDC} k')^{-1}$$

b_{IDC} : b coefficients for IDC¹

Σ_{IDC} : symmetrical matrices for IDC¹

K : Pure spectra of interest (peanut)




Fig b. IDC loadings (b coefficient)

2. Loadings application on HS images

2.1 HS images pretreatment: SNV and SAVGOL

2.2 Application of the b coefficients to obtain score images (Fig. d.)




Fig c. Score validation for pure

3. Image segmentation

3.1 Score images were segmented into flour and peanut pixels according to a reference image histogram.

Results




Fig d. Scores image Fig e. Class image

- Classification images provided clear detection of peanut traces in wheat flour Fig.(e).
- Minimum level of peanut traces detected with present experimental setup was 0.01 % by weight.

Conclusions

- NIR Hyperspectral imaging (1000-2500 nm) combined with IDC allowed the detection of peanut traces down to adulteration percentages 0.01%.
- Contrary to PLSR, IDC does not require a calibration set, but uses both expert and experimental information and suitable for quantification of an interest compound in complex matrices.
- The obtained results shows the feasibility of using HSI systems for the detection of peanut traces in conjunction with chemical procedures, such as RT-PCR and ELISA.

References

1. Boulet, J. & Roger, J. 2010, "Improvement of Direct Calibration in spectroscopy", *Analytica Chimica Acta*, vol. 668, no. 2, pp. 130-136.
2. Cochard, M.M. & Eigenmann, P.A. 2011, "Chapter 15 - Allergies to Nuts and Seeds" in *Nuts and Seeds in Health and Disease Prevention*, eds. V.R. Preedy, R.R. Watson & V.B. Patel, Academic Press, San Diego, pp. 137-143.
3. Hostetler, T.L., Hostetler, S.G., Phillips, G. & Martin, B.L. 2012, "The ability of adults and children to visually identify peanuts and tree nuts", *Annals of Allergy, Asthma & Immunology*, vol. 108, no. 1, pp. 25-29.

The authors are grateful for peanut sample sets used in the work provided by European Commission Joint Research Centre, Institute for Reference Materials and Measurements and funding to attend the conference provided under Erasmus Mundus INDIA4EUll program and Campus of International Excellence, Moncloa.